

Note on the Sunyaev-Zel'dovich thermal effect

A. Sandoval-Villalazo^a and L.S. García-Colín^{b, c}

^a Departamento de Física y Matemáticas, Universidad Iberoamericana

Lomas de Santa Fe 01210 México D.F., México

E-Mail: alfredo.sandoval@uia.mx

^b Departamento de Física, Universidad Autónoma Metropolitana

México D.F., 09340 México

^c El Colegio Nacional, Centro Histórico 06020

México D.F., México

E-Mail: lgcs@xanum.uam.mx

February 2, 2008

Abstract

In a previous publication we have derived an expression for the full distorted spectrum arising when the photons of the cosmic background radiation are absorbed and emitted by an optically thin gas. The expression simply adds up the effects of the joint probability that an electron absorbs an incoming photon of frequency $\bar{\nu}$ and emits it with a frequency ν . We here show that such an expression is identical to the Kompaneets equation describing the thermal Sunyaev-Zel'dovich effect. This clearly shows that the two pictures, diffusion and absorption-emission are mathematically equivalent to first order in the Compton parameter. Other consequences of this result are also discussed.

In a recent publication [1] we have given a detailed discussion of how the thermal Sunyaev-Zel'dovich (SZ) effect [2] [3] may be interpreted in terms of a rather simplified picture as an absorption-emission process. In fact, our view is that such an effect is indeed that of an absorption-emission process in which a few photons happen to be captured by electrons in the optically thin gas. An electron moving with a given thermal velocity emits (scatters) a photon with a certain incoming frequency $\bar{\nu}$ and outgoing frequency ν . The line breath of this process is readily calculated from kinetic theory taking into account that the media in which the process takes place has a small optical depth directly related to the Compton parameter y . When the resulting expression, which will be called *structure factor*, is convoluted with the incoming flux of photons obtained from Planck's distribution, one easily obtains the disturbed spectra.

In Ref. [1] we derived an analytical expression for the full distorted spectrum

of the scattered radiation which reads as

$$I(\nu) = \frac{1}{\sqrt{\pi}W(\nu)} \int_0^\infty I_o(\bar{\nu}) \exp \left[- \left(\frac{\bar{\nu} - (1-2y)\nu}{W(\nu)} \right)^2 \right] d\bar{\nu} \quad (1)$$

and defines the joint probability of finding an electron scattering a photon with incoming frequency $\bar{\nu}$ and outgoing frequency ν multiplied by the incoming photon flux $I_o(\bar{\nu})$. In Eq. (1) the incoming flux is given by

$$I_o(\bar{\nu}) = \frac{2h\bar{\nu}^3}{c^2} \frac{1}{e^{\frac{h\bar{\nu}}{kT}} - 1} \quad (2)$$

where $T \equiv 2.726 \text{ K}$, the temperature of the cosmic background radiation, k is Boltzmann's constant, h is Planck's constant and c the speed of light. Further, $W(\nu)$ is the width of the spectral line at frequency ν and its squared value reads (Eq. (14) in Ref. [1]):

$$W^2(\nu) = \frac{4kT_e}{m_e c^2} \tau \nu^2 = 4y\nu^2 \quad (3)$$

where T_e is the temperature of the electron scatterers, m_e is the electron mass and τ is the optical depth, whose meaning is fully discussed in [1].

The purpose of this note is simply to show that Eq. (1) can be evaluated analytically in an almost trivial way leading to an expression which is identical to the one reported in the literature. Setting in Eq. (1)

$$\alpha = \frac{\bar{\nu} - (1-2y)\nu}{W(\nu)} \quad (4)$$

we get that

$$I(\nu) = \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} I_o(\nu + \Delta\nu) \exp[-\alpha^2] d\alpha \quad (5)$$

where

$$\Delta\nu = -2y\nu + 2y^{1/2}\nu\alpha \quad (6)$$

Notice should be made on the fact that the lower limit in the integral can be readily estimated to be a large negative number, which is for all practical purposes $-\infty$. The trick is now immediate. Since y is a very small number [4] and I_o is an analytical function of ν we may expand the integrand in Eq. (5) in a Taylor series. Since $\Delta\nu$ is linear in α , all odd powers of α yield zero upon integration so that after a straightforward calculation we get that

$$I(\nu) - I_o(\nu) = -2y\nu \frac{\partial I_o}{\partial \nu} + y\nu^2 \frac{\partial^2 I_o}{\partial \nu^2} + 2y^2\nu^2 \frac{\partial^2 I_o}{\partial \nu^2} - 2y^2\nu^3 \frac{\partial^3 I_o}{\partial \nu^3} + \dots \quad (7)$$

To first order in y we reproduce *exactly* the analytic expression for the distortion curve [2]-[3]. It is also interesting to notice that, taking into account the second

order terms in y included in Eq. (7), the crossover frequency ν_c corresponding to the solution of equation

$$I(\nu) - I_o(\nu) = 0 \quad (8)$$

is now clearly dependent on the value of y namely, on the electron temperature, a feature not ordinarily recognized for a non-relativistic intra-cluster gas. For a typical value of $y = 10^{-4}$, we obtain $\nu_c = 217.16$ GHz, while the value obtained by considering only linear terms in y yields $\nu_c = 217.12$ GHz, a difference of nearly 40 MHz.

The result expressed in Eq. (7) is of significant importance. Not only shows that laborious calculations may be expressed in terms of rather elementary concepts borrowed from statistical mechanics, but clarifies an until now intriguing result. Indeed, Eq. (7) contains the famous Kompaneets equation [5]-[7] describing the thermal SZ effect using a diffusive mechanism to account for the motion of photons in an optically thin electron gas where the electron temperature $T_e \gg T_{CMB}$, the photon temperature (see specifically Ref. [6]). One must notice that, since most photons *are not scattered even once*, a diffusion approximation would then hardly seem adequate [4]. Nevertheless, the diffusion mechanism is mathematically identical to the absorption-emission process of photons by electrons to first order in y . The problem posed in Ref. [1] requiring an explanation of why two different mechanisms lead to the same result is now mathematically solved. The exponential function in the integrand of Eq.(1) is playing the role of a Green's function for the diffusion equation. Finally, we want to stress that it is now clear why the curves describing the thermal SZ effect obtained by numerical integration of different mathematical expressions turns to be identical. We therefore expect that the relativistic SZ effect and other photon scattering problems in hot plasmas can be handled by a similar procedure.

The authors wish to thank G.H. Weiss for kindly supplying the idea behind the integration method. This work has been supported by CONACyT (Mexico), project 41081-F.

References

- [1] A. Sandoval-Villalbazo and L.S. García-Colín; J.Phys. A: Math and Gen. **36** (2003) 4641-4650 [astro-ph/0208440].
- [2] Y.B. Zel'dovich and R.A. Sunyaev, Astrophys. Space Sci. **4**, 301 (1969).
- [3] R.A. Sunyaev and Ya. B. Zel'dovich. Comm. Astrophys. Space Phys. **4**, 173 (1972).
- [4] Y. Rephaeli, Astrophys. J. **445**, 33 (1995);
A.D. Dolgov, S.H. Hansen, D.V. Semikoz and S. Pastor, Astrophys. J. **554** 74 (2001) [astro-ph/0010412].

- [5] P.J.E. Peebles, Physical Cosmology (Princeton Univ. Press, Princeton, 2nd. Ed. 1993).
- [6] R.A. Sunyaev and Ya. B. Zel'dovich; Ann. Rev. Astrom. Astrophys. **18**, 537 (1980)
- [7] A.S. Kompaneets; JETP. **4**, 730 (1957).